

QCD PHASE SHIFTS AND RISING TOTAL CROSS SECTIONS

C.S. Lam

Department of Physics, McGill University, 3600 University St., Montreal, QC, Canada H3A 2T8
E-mail: Lam@physics.mcgill.ca

Energy dependence of γ^*p total cross section is considered. It is known from the HERA data that the cross section grows with energy, and the rate of growth is an increasing function of the virtuality Q^2 of the photon γ^* . This dependence can be explained in a simple and generic way by the Froissart bound. To implement this mechanism quantitatively a theory satisfying s -channel unitarity is required. This is achieved by computing the total cross section from the optical theorem, and the forward elastic amplitude from the phase shifts in the impact-parameter representation. A recipe to do so in perturbative QCD is discussed, together with an expose of the advantage to calculate the elastic amplitude via phase shift, rather than direct sum of all Feynman diagrams. A two-loop computation of phase shift and total cross section is presented, and compared with the HERA data with good agreements.

1 Total Cross Sections

1.1 Experimental Data

γ^*p total cross section $\sigma_{tot}(s, Q)$ measured at HERA^{1,2} can be fitted by the formula $\sigma_0(Q)(s/\Lambda^2)^{a(Q)}$, where s is the square of the c.m. energy, and Q^2 is the virtuality of the photon. The power $a(Q)$ is equal to 0.08 at $Q = 0$, the same as the corresponding power for *all* hadronic total cross sections³. We shall refer to this experimental feature, that the energy dependence of the total cross section is independent of the nature of the colliding particles, as *universality*. Moreover, $a(Q)$ is an increasing function of Q . Asymptotically, total cross sections are limited by the Froissart bound not to grow faster than $\ln^2 s$, so the power-like increase observed must be approximate, with the effective power of growth decreasing to zero at infinite energies.

1.2 Q -Dependence of $a(Q)$

This last remark provides a simple and generic mechanism to explain why $a(Q)$ is an increasing function of Q . From universality, we can write $\sigma_{tot}(s, Q)$ in the form

$$\sigma_{tot}(s, Q) = \sigma_0(Q)f(s'), \quad (1)$$

where $\sigma_0(Q)$ depends both on Q and the nature of the colliding particles, but $f(s')$, which governs the energy dependence of the total cross sections, is universal and independent of the nature of the colliding particles. It is a dimensionless function of the dimensionless energy $s' = s/\Lambda^2(Q)$. On dimensional grounds, the energy-scale parameter $\Lambda(Q)$ is expected to increase with Q .

To satisfy the Froissart bound, the universal function $f(s')$ must asymptotically approach $\ln^2 s'$. This means that at sufficiently high energies, its slope in a $\ln f(s')$ vs $\ln s'$ plot must increase with decreasing s' . A larger Q leads to a larger $\Lambda(Q)$, a smaller s' , and hence a larger slope $a(Q)$ in the log-log plot. This shows that $a(Q)$ is an

increasing function of Q , just as observed in the HERA data^{1,2}.

2 QCD Phase Shifts

2.1 Impact-Parameter Representation

To be able to implement this mechanism quantitatively, we must start from a theory capable of describing rising total cross sections which obey the Froissart bound. This is so if total cross sections are computed from the optical theorem, and the forward elastic scattering amplitudes are calculated from phase shifts in the impact-parameter representation:

$$\sigma_{tot}(s) = \frac{1}{s} \text{Im} A_0(s, 0) \quad (2)$$

$$A(s, \Delta) = -is \int d^2b e^{i\vec{\Delta} \cdot \vec{b}} \left(e^{2i\delta(s, b)} - 1 \right) \quad (3)$$

The impact parameter $\vec{b}$ and its conjugate, the momentum transfer $\vec{\Delta}$, are two-dimensional vectors in the transverse plane. In QCD, the amplitude A is a colour matrix whose matrix elements are the scattering amplitudes of partons of various colours. The amplitude A_0 appearing in (2) is obtained from A of (3) by retaining only the colour-singlet exchange amplitude.

The effective size R measured in the total cross section can be estimated from the condition $\delta(s, R) \sim 1$. If the interaction has a range μ^{-1} , so that $\delta(s, b) \sim \xi(s) \exp(-\mu b)$ for large b , then the effective radius is given by $R = \mu^{-1} \ln \xi(s)$. This grows with energy if $\xi(s)$ does, and as long as $\xi(s)$ does not grow faster than a power, R will not rise more rapidly than $\ln s$, and $\sigma_{tot} \sim \pi R^2$ will not grow faster than $\ln^2 s$, as prescribed by the Froissart bound.

2.2 QCD Phase Shifts

So far the discussion has been general and qualitative. To proceed further we must find a way to calculate the

phase shift $\delta(s, b)$. We shall discuss how to do so within the framework of perturbative QCD.

Scattering amplitudes are given by Feynman diagrams. From (3), the phase shift is given by the logarithm of the amplitude in the impact-parameter space. With the presence of non-commuting colour matrices in the amplitude this calculation promises to be complicated. It is not a priori clear that any simple formula can be obtained for the phase shift. Fortunately, in the leading-log approximation (LLA) where only the highest power of $\ln s$ is kept, the result turns out to be very simple^{4,5,6,7,8,9}. We will now state this result for quark-quark scattering.

First, use the $SU(N_c)$ commutation relation $[\lambda_a, \lambda_b] = if_{abc}\lambda_c$ for colour matrices λ_a in any representation, together with the normalization condition $\text{Tr}(\lambda_a\lambda_b) = \delta_{ab}/2$ for the colour matrices in the fundamental representation, to reduce the colour factor of each Feynman diagram into combination of a linearly independent set of ‘pseudo-planar’ colour factors, as illustrated in Fig. 1. Choose from this set all the *primitive colour factors* G_i , which are those that remain connected when the two quark lines are removed from the diagram. Of those shown in Fig. 1, G_a and G_d are primitive, and G_b, G_c, G_e, G_f are not. Since many Feynman diagrams can contribute to the same primitive colour factor G_i , the corresponding amplitude $d_i(s, b)$ in the impact-parameter space will generally be given by a sum of many Feynman diagrams.

The phase shift is obtained by summing the contributions from all primitive colour diagrams:

$$2\delta(s, b) = \sum_i d_i(s, b)G_i. \quad (4)$$

It is clear from (3) that a $\delta(s, b)$ calculate even in the lowest order will give rise to an expression for the amplitude $A(s, \Delta)$ involving amplitudes of all orders. In this way a phase shift calculation of an amplitude is similar to a renormalization-group improved calculation of a Green’s function or a physical parameter. Moreover, it can also be shown⁶ that the g^2 and $\ln s$ dependences of $d_i(s, b)$ are of the form $g^{2m}D(g^2 \ln s)$, if G_i has m gluon lines connecting the two quarks ($m = 1, 2, 3, 2, 4, 4$ respectively for (a), (b), (c), (d), (e) in Fig. 1). Let us illustrate the significance of these two remarks.

Suppose we calculate electron-electron scattering in QED, via the exchange of n photons. We will ignore all fermion loops in the intermediate states. There are $n!$ (crossed and uncrossed) ladder diagrams describing this process, whose individual powers of $\ln s$ grow with n . However, it is known that when the $n!$ diagrams are summed, *all* powers of $\ln s$ cancel one other, leaving behind a contribution to the total cross section independent of energy. This can be seen from (4) in the following




Figure 1: Colour diagrams. The colour factors (a) and (d) are primitive, the others are not.

way. In this case, the only primitive ‘colour’ diagram is a single-photon exchange diagram, G_a , for all the others are disconnected once the two electron lines are removed (G_d is not present for QED). Among the $n!$ ladder diagrams, the only one that would contribute to G_a is the Born amplitude, with $n = 1$. Hence the phase shift is given by the Born amplitude, which is proportional to g^2 and contains no $\ln s$. This agrees with the general form discussed above with $m = 1$. Moreover, when we expand the exponential in (3) to order g^{2n} , we see that the amplitude to that order must also contain no $\ln s$ terms, agreeing with the result obtained by direct calculations.

For quark-quark scattering in QCD, gluon self coupling allows an infinite number of primitive colour factors to be formed, even when quark loops are omitted. G_a is colourwise a single gluon exchange, with amplitude of the form $d_a(s, b) \sim g^2 D(g^2 \ln s)$. When all diagrams are summed in LLA, this gives rise to the familiar Reggeized gluon amplitude with D being an exponential function. Otherwise D is a polynomial if the sum is taken only to a finite order. When the exponential in (3) is expanded with $\delta(s, b) \sim d_a G_a$ being the exponent, the constant term is cancelled and the linear term has colour-octet exchange, so it is starting from the quadratic term that it may contribute to the colour-singlet amplitude A_0 in (2). For that reason the contribution from G_a to σ_{tot} , is of order g^4 , times a function of $g^2 \ln s$. In the region where g^2 is small and $g^2 \ln s$ is of order unity, the contribution from G_a to σ_{tot} is then $O(g^4)$. The only other terms of the same order came from primitive colour factors that have two vertical gluon lines between the two quarks ($m = 2$), such as G_d in Fig. 1. Those diagrams are also of $O(g^4)$, and since two vertical gluons are involved, the primitive colour factors do contribute directly to a colour-singlet exchange. The BFKL Pomeron amplitude is therefore the $O(g^4)$ contributions from the $m = 1$ and the $m = 2$ primitive colour factors. To be sure, other terms in the expansions of the exponential give higher order contributions, but these subleading-log terms are necessary to uphold s -channel unitarity and the Froissart bound.

3 Two-Loop Phase Shifts

The quark-quark scattering amplitude in LLA, up to two loops or $O(g^6)$, can be found in the book of Cheng and Wu¹⁰. All we need to do is to extract from it the phase shift $\delta(s, b)$ using eq. (4), and then the colour-singlet exchange amplitude A_0 in (2). The result for the universal energy function $f(s')$ in (1) is given by⁸

$$f(s') = \int d^2x \left[1 - \frac{2}{3} e^{-D} \cos\left(\frac{2\delta_a}{3}\right) - \frac{1}{3} e^{-2D} \cos\left(\frac{4\delta_a}{3}\right) \right]$$

$$2\delta_a(s, b) = -g^2 K_0(x) + (g^4 N_c \ln s' / 4\pi) K_0^2(x) - (g^6 N_c^2 \ln^2 s' / 32\pi^2) V(x),$$

$$D(x) = -i\delta_d(x) = (g^6 \ln s' / 4\pi) [K_0^3(x) - \frac{1}{2} V(x)], \quad (5)$$

where $K_0(x)$ is the Bessel K -function, and $V(x) = -(1/2\pi)^4 \nabla_x^2 \int d^2x' K_0^2(|\vec{x} - \vec{x}'|) K_0^2(|\vec{x}'|)$.

4 Comparison with the HERA Data

Equation (5) is computed for quark-quark scattering, but if the energy function $f(s')$ is indeed universal, as experimental data indicate, (see the discussion leading to (1)), then we can use it to compare with experimental results. Even so, the perturbative formula in (5) is expected to be quantitatively accurate only when both g^2 and $g^2 \ln s'$ are small. The former is so when the virtuality Q^2 is large, but the latter is not satisfied for the present data. Higher-order calculations are needed and they are underway⁹, but for now, to get a qualitative idea, let us compare (5) with data anyhow. Choosing a linear dependence for the energy scale parameter, $\Lambda(Q) = \Lambda_0 + cQ$, with $\Lambda_0 = 0.2$ GeV and $c = 4$, the comparison is shown in Fig. 1. We see that inspite of the small Q available in the data, the agreement of theory and data seems to be very good.

This agreement can probably be attributed to two things. First, the mechanism explained in section 1.2 is so general that any reasonable function $f(s')$ satisfying the Froissart bound is expected to yield a reasonable result, if a suitable energy-scale parameter $\Lambda(Q)$ is chosen. Moreover, the agreement is reminiscent of the success in the early days of QCD, before higher-loop formulas became available. Drell-Yan pair mass spectrum and other quantities were correctly predicted, though normalizations were off by the famous ‘K factor’. When higher-order corrections are taken into account, there will be a correction to the universal energy function $f(s')$ with a corresponding change in the energy-scale parameter $\Lambda(Q)$ to fit the data. This is not unlike Λ_{QCD} , which takes on different values when deduced from experimentally measured $\alpha_s(Q)$, depending on whether it appears in a one-loop or a two-loop formula.




Figure 2: γ^*p total cross-sections as a function of s . Data are taken from Ref. ¹ where references to the original experiments can be found. Q^2 of the photon, in $(\text{GeV})^2$, are given on the right. Where two numbers are listed, the second one refers to the low-energy data of E665. The solid curve is the prediction of the two-loop phase shift calculation, eq. (5). The dotted lines are $s^{0.08}$, and the dash line is $s^{0.05}$, both there for comparisons.

5 Conclusion

Experimentally γ^*p total cross section grows with energy like $s^{a(Q)}$, with an effective power $a(Q)$ increasing with the virtuality Q^2 of the photon γ^* . This behaviour can be explained by universality and the Froissart bound. To compute the universal energy function $f(s')$ governing the energy dependence of all total cross sections, we resort to the impact-parameter formalism which preserves s -channel unitarity and the Froissart bound. We have discussed a recipe in perturbative QCD to calculate the phase shift $\delta(s, b)$ needed in that formalism. We have also discussed the advantage of calculating the elastic amplitude using phase shift, rather than doing it directly from summing all Feynman diagrams. Formulas are given for quark-quark phase shifts to two loops, and for the corresponding energy function $f(s')$. When they are used to compare with the HERA experimental data, good agreements are obtained, the reasons for which are also dis-

cussed.

References

1. ZEUS Collaboration, hep-ex/9707025.
2. H1 Collaboration, hep-ex/9708016; ZEUS Collaboration, hep-ex/9709021, 9602010.
3. A. Donnachie and P.V. Landshoff, *Phys. Lett.* **B296** (1992) 227.
4. C.S. Lam and K.F. Liu, *Nucl. Phys.* **B483** (1997) 514.
5. Y.J. Feng, O. Hamidi-Ravari, and C.S. Lam, *Phys. Rev. D* **54** (1996) 3114.
6. Y.J. Feng and C.S. Lam, *Phys. Rev. D* **55** (1997) 4016.
7. C.S. Lam, hep-th/9804181, to appear in the *Journal of Mathematical Physics*.
8. C.S. Lam, hep-ph/9804463.
9. R. Dib, J. Khoury, and C.S. Lam, to be published.
10. H. Cheng and T.T. Wu, *'Expanding Protons: Scattering at High Energies'*, (M.I.T. Press, 1987).